Digitized by the Internet Archive
in 2026

https://archive.org/details/journal-of-chemical-thermodynamics_1989_21_contents

THE JOURNAL OF CHEMICAL THERMODYNAMICS

Volume 21, 1989

EDITORS

M. L. McGlashan
Department of Chemistry
University College London
20 Gordon Street
London WC1H 0AJ
U.K.

P. A. G. O'Hare
National Institute of
Standards and Technology
Building 222, Gaithersburg
MD 20899
U.S.A.

Assistant Editor: **M. B. Ewing** (University College London, U.K.)

ADVISORY BOARD

Chairman: G. M. Schneider (F.R.G.)

J. C. G. Calado (Portugal)
H. Chihara (Japan)
E. H. P. Cordfunke (The Netherlands)
J. H. Dymond (U.K.)
J. Fuger (Commission of the European Communities)
M. Gaune-Escard (France)
L. V. Gurvich (U.S.S.R.)
C. Jolicoeur (Canada)
K. N. Marsh (U.S.A)
G. Pilcher (U K.)
G. Somsen (The Netherlands)
W. V. Steele (U.S.A.)
F. L. Swinton (U.K.)
W. Wagner (F.R.G.)
R. D. Weir (Canada)
R. H. Wood (U.S.A.)
W. A. Wóycicki (Poland)

ACADEMIC PRESS
(Harcourt Brace Jovanovich, Publishers)
London San Diego New York
Boston Sydney Tokyo Toronto

30
D
21
683

Copyright © 1989 Academic Press Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 21

Number 1, January 1989

PITZER, K. S. Fluids, both ionic and non-ionic, over wide ranges of temperature and composition. (The seventh Rossini Lecture of the Commission on Thermodynamics of IUPAC delivered on 29 August 1988 at its Conference on Chemical Thermodynamics held in Prague, Czechoslovakia) . . 1

LETCHER, T. M.; SEWRY, J.; BAXTER, R. Excess molar volumes of (a cycloalkane + an alk-1-yne) at 298.15 K 19

FROLOVA, G. I.; REZNIK, L. E.; PAUKOV, I. E. Crystal-field effects on the thermodynamic properties of the $KLn(MoO_4)_2$ isostructural compounds. 25

BUGAJEWSKI, Z.; BYLICKI, A. (Solid + liquid) phase equilibria for (*N*-methylpyrrolidin-2-one + *p*-cresol) 37

PANDO, C.; RENUNCIO, J. A. R.; FAUX, P. W.; CHRISTENSEN, J. J.; IZATT, R. M. The excess enthalpies of (carbon dioxide + *n*-hexane + toluene) at 573.15 K and 7.50 and 12.50 MPa 41

ERNST, G.; MAURER, G.; WIEDERUH, E. Flow calorimeter for the accurate determination of the isobaric heat capacity at high pressures; results for carbon dioxide. 53

WANG, L.; BENSON, G. C.; LU, B. C.-Y. Excess enthalpies and excess volumes for (di-*n*-propyl ether + a cycloalkane) 67

MORIYOSHI, T.; NAKAGAWA, M. Compressions of aqueous binary mixtures containing amides and carbamic-acid derivatives at 298.15 K and 101.3 MPa 73

STEELE, W. V.; CHIRICO, R. D.; HOSSENLOPP, I. A.; NGUYEN, A.; SMITH, N. K.; GAMMON, B. E. The thermodynamic properties of the five benzoquinolines. 81

ADVICE TO AUTHORS 109

Number 2, February 1989

LOOS, TH. W. DE; POOT, W.; SWAAN ARONS, J. DE. Fluid-phase equilibria and critical phenomena in (tetrafluoromethane + *n*-butane) 113

RAI, R. D.; SHUKLA, R. K.; SHUKLA, A. K.; PANDEY, J. D. Ultrasonic speeds and isentropic compressibilities of ternary liquid mixtures at (298.15 ± 0.01) K. 125

RAMACHANDRAN, T. P.; RAMACHANDRAN, S.; ANANTH, M. S. Isothermal (vapour + liquid) equilibria of three binary mixtures at four temperatures. 131

VIŠIĆ, M.; MEKJAVIĆ, I. Thermodynamics of the cell Cd(s) + Hg|$CdCl_2$ (aq, *m*) |AgCl|Ag 139

WANG, L.; BENSON, G. C.; LU, B. C.-Y. Excess volumes for (di-*n*-propyl ether + *n*-alkane) at 298.15 K 147

O'HARE, P. A. G.; AWERE, E. G.; PARSONS, S.; PASSMORE, J. The standard molar enthalpy of formation at 298.15 K of $S_2N^+AsF_6^-$ by fluorine-combustion calorimetry 153

RIBEIRO DA SILVA, M. A. V.; MONTE, M. J. S.; MATOS, M. A. R. Enthalpies of combustion, vapour pressures, and enthalpies of sublimation of 8-hydroxyquinoline, 5-nitro-8-hydroxyquinoline, and 2-methyl-8-hydroxyquinoline . 159

RIBEIRO DA SILVA, M. A. V.; REIS, A. M. M. V. Enthalpies of formation of five crystalline dialkyldithiocarbamate copper(II) complexes. The mean Cu–S bond-dissociation enthalpies 167

RIBEIRO DA SILVA, M. D. M. C.; SOUZA, P.; PILCHER, G. Enthalpies of combustion of thiobenzamide, *N*,*N*-dimethylthiobenzamide, and *N*,*N*-diethylthiobenzamide 173

CHIRICO, R. D.; HOSSENLOPP, I. A.; NGUYEN, A.; STEELE, W. V.; GAMMON, B. E. The thermodynamic properties of 4-methylphenanthrene 179

KNAUTH, P.; SABBAH, R. Combustion calorimetry on milligram samples of liquid substances with a CRMT rocking bomb calorimeter. Application to the study of ω-alkanediols at 298.15 K 203

MARSH, K. N.; ROGERS, H. Excess enthalpies of (cyclohexane + nitroethane) near the (liquid + liquid) critical state 211

MORIYOSHI, T.; UOSAKI, Y.; SAKAMOTO, T.; HAYASHI, Y. (Liquid + liquid) equilibria of (water + ethanol + 2,6-dimethyl-4-heptanone) from 0.1 to 200 MPa at 298.15 and 323.15 K 219

Number 3, March 1989

NAGATA, I. Isothermal (vapour + liquid) equilibria of (2-methylpropan-2-ol) + acetonitrile) and (2-methylpropan-2-ol + acetonitrile + benzene) . . 225

GARCIA BAONZA, V.; NUÑEZ DELGADO, J.; CACERES ALONSO, M. (p, ρ, T) properties of liquid carbon disulfide from 198 to 298 K and pressures up to 104 MPa 231

IMAMURA, A.; TAKAHASHI, K.; MURATA, S.; SAKIYAMA, M. Standard enthalpies of formation of trimethyl cyanurate, malonamide, and 1,3-dimethyluracil. 237

RUDNYI, E. B.; VOVK, O. M.; KAIBICHEVA, E. A.; SIDOROV, L. N. Formation enthalpies of oxygen-containing anions of Group VI elements in the gas phase and the electron affinities of CrO_3, MoO_3, and WO_3 . . . 247

PÉREZ, P.; VALERO, J.; GRACIA, M.; GUTIÉRREZ LOSA, C. G^E_m(298.15 K) of mixtures containing 1,2-dichloropropane or 1,3-dichloropropane . . 259

RIBEIRO DA SILVA, M. A. V.; RIBEIRO DA SILVA, M. D. M. C.; TEIXEIRA, J. A. S.; BRUCE, J. M.; GUYAN, P. M.; PILCHER, G. Enthalpies of combustion of 1,4-naphthoquinone, 9,10-anthraquinone, 9,10-phenanthraquinone, 1,4,9,10-anthradiquinone, 5,8-dihydroxy-1,4-naphthoquinone, and 1,4-dihydroxy-9,10-anthraquinone 265

COLOMINA, M.; JIMÉNEZ, P.; ROUX, M. V.; TURRIÓN, C. Thermochemical properties of 1,2,4,5-tetramethylbenzene, pentamethylbenzene, and hexamethylbenzene 275

AIROLDI, C.; SOUZA, A. G. DE. Thermochemical properties of chelates of di-*n*-butyldithiocarbamate with phosphorus-group elements . . . 283

NICKEL, D.; SCHNEIDER, G. M. Near-infrared spectroscopic investigations on phase behaviour and association of 1-hexanol and 1-decanol in carbon dioxide, chlorotrifluoromethane, and sulfur hexafluoride . . . 293

FRIEDRICH, J.; SCHNEIDER, G. M. Near-infrared spectroscopic investigations on phase behaviour and association of 1-octadecanol, 3-hexanol, and 3-methyl-3-pentanol in carbon dioxide and chlorotrifluoromethane . 307

MAJER, V.; INGLESE, A.; WOOD, R. H. Volumetric properties of LiCl(aq) from 0.05 to 3.0 mol·kg^{-1}, 322 to 550 K, and 0.8 to 32.6 MPa. . . . 321

SHARMA, S. C.; JOSHI, I. M.; SINGH, J. Excess molar volumes of (oxolane + trichloromethane or tetrachloromethane or 1,2-dichloroethane or 1,1,2-trichloroethane or 1,1,2,2-tetrachloroethane or 1,1,1,2,2-pentachloroethane) at 303.15 K 331

Number 4, April 1989

CHAO, J. P.; DAI, M. Studies on thermodynamic properties of binary mixtures containing an alcohol. XI. Excess enthalpies of each of (one of the four butanols + chlorobenzene or aniline) 337

SIJPKES, A. H.; OUDHUIS, A. A. C. M.; SOMSEN, G.; LILLEY, T. H. Enthalpies of solution of amides and peptides in aqueous solutions of urea and in *N*,*N*-dimethylformamide at 298.15 K 343

HOLMES, R. D.; KERSTING, A. B.; ARCULUS, R. J. Standard molar Gibbs free energy of formation for Cu_2O: high-resolution electrochemical measurements from 900 to 1300 K 351

Gonzalez-Alvarez, D.; Grønvold, F.; Falk, B.; Westrum, E. F., Jr.; Blachnik, R.; Kudermann, G. FeAs: Heat capacity, enthalpy increments, other thermodynamic properties from 5 to 1350 K, and magnetic transition 363

Labban, A. K.; López de la Fuente, F. L.; Cheda, J. A. R.; Westrum, E. F., Jr.; Fernández-Martín, F. Thermodynamics of thallium alkanoates. VI. Thallium(I) *n*-heptanoate revisited 375

Nesterova, T. N.; Pimerzin, A. A.; Rozhnov, A. M.; Karlina, T. N. Equilibria for the isomerization of (secondary-alkyl)phenols and cyclohexylphenols . 385

Majer, V.; Inglese, A.; Wood, R. H. Volumetric properties of NaBr(aq, $m = 0.05$ to 3.0 mol·kg^{-1}, $T = 322$ to 550 K, $p = 0.8$ to 32.6 MPa) 397

Ernst, G.; Hochberg, U. E. Flow-calorimetric results for the specific heat capacity c_p of CO_2, of C_2H_6, and of $(0.5CO_2 + 0.5C_2H_6)$ at high pressures. 407

Konings, R. J. M.; Cordfunke, E. H. P.; Ouweltjes, W. The standard enthalpies of formation of hydroxides. III. LiOH 415

Fernández, J.; Velasco, I.; Otín, S. Excess molar enthalpies of (cyclohexane + an *n*-alkylamine) 419

Ribeiro da Silva, M. A. V.; Reis, A. M. M. V. Enthalpies of formation of five crystalline dialkyldithiocarbamate nickel(II) complexes: the mean (Ni–S) bond-dissociation enthalpies 423

Escarda, R.; Garcia, I.; Cobos, J. C.; Casanova, C. Excess molar volumes of (butanoic anhydride or heptanoic anhydride + *n*-alkane) at 298.15 K . 431

Markó, L.; Jákli, Gy.; Jancsó, G. Vapour pressure of heavy water at its triple point 437

Acree, W. E., Jr.; Kirchner, J. J.; Tucker, S. A.; Pilcher, G.; Ribeiro da Silva, M. D. M. C. Enthalpies of combustion of three benzylidene *t*-butylamine *N*-oxide derivatives and of 4-nitrobenzylidene *t*-butylamine: the dissociation enthalpies of the (N–O) bonds 443

Number 5, May 1989

Kirklin, D. R.; Domalski, E. S. Energy of combustion of triphenylphosphate. 449

Nagata, I. (Liquid + liquid) equilibria of (acetonitrile + 2-butanol or 2-methyl-2-propanol + cyclohexane or *n*-hexane or *n*-heptane) . . 457

Rard, J. A.; Miller, D. G. Isopiestic determination of the osmotic and activity coefficients of $ZnCl_2$(aq) at 298.15 K 463

FERRO, D.; PIACENTE, V.; SCARDALA, P. Decomposition enthalpies of iron sulfides 483

NÚÑEZ, L.; MIGUELEZ, F.; BARRAL, L.; PAZ ANDRADE, M. I. Excess molar enthalpies at 298.15 K of (1-chloropentane + nonane or decane) and of (1-chlorohexane + nonane or decane or undecane) 495

MAYRATH, J. E.; MAGEE, J. W. Measurements of molar heat capacity at constant volume: $C_{V,m}\{xCH_4+(1-x)C_2H_6,\ T = 100$ to 320 K, $p \leqslant 35$ MPa$\}$ 499

ALONSO, R.; CORRALES, J. A. Excess molar volumes of (methylcyclohexane + an alkanol) at 298.15 K. I. Results for *n*-butanol, *n*-pentanol, and *n*-hexanol 515

HALLÉN, D.; WADSÖ, I. A new microcalorimetric vessel for dissolution of slightly soluble gases. Enthalpies of solution in water of carbon tetrafluoride and sulphur hexafluoride at 288.15, 298.15, and 308.15 K . 519

HALLÉN, D.; NILSSON, S.-O.; WADSÖ, I. A new flow-microcalorimetric vessel for dissolution of small quantities of easily or slightly soluble liquids . . 529

RARD, J. A. Isopiestic determination of the osmotic and activity coefficients of $\{(1-y)H_2SO_4 + yNa_2SO_4\}$(aq) at 298.15 K. I. Results for $y = 0.5$ ($NaHSO_4$) and $y = 0.55595$, 0.70189, and 0.84920 539

Number 6, June 1989

SIMONSON, J. M.; MESMER, R. E.; ROGERS, P. S. Z. The enthalpy of dilution and apparent molar heat capacity of NaOH(aq) to 523 K and 40 MPa . . 561

SACCHETTO, G. A.; KODEJŠ, Z. Vapour pressures of very concentrated electrolyte solutions. V. Measurements on $\{(1-x)H_2O + xKSCN\}$ by a dew-point apparatus 585

ROGERS, P. S. Z.; DUFFY, C. J. Comparison of calibration methods for flow heat-capacity calorimeters and heat capacities of concentrated NaCl(aq) to 598 K 595

TAN, Z. C.; KAMAGUCHI, A.; NAGANO, Y.; SAKIYAMA, M. Standard enthalpy of formation of 2-chloro-6-(trichloromethyl)pyridine 615

MUSSINI, P. R.; LONGHI, P.; MUSSINI, T.; RONDININI, S. The second ionization constant of aqueous sulphuric acid at 298.15 K from the electromotive force of the unbuffered cell: $H_2(g)|H_2SO_4(aq)|Hg_2SO_4(s)|Hg$ . . . 625

SHAVIV, R.; WESTRUM, E. F., JR.; GRØNVOLD, F.; STØLEN, S.; INABA, A.; FUJII, H.; CHIHARA, H. Heat capacity, thermodynamic properties, and transitions of silver iodide 631

WANG, Z.-C.; ZHANG, X.-H.; HE, T.-Z.; BAO, Y.-H. High temperature isopiestic studies on $\{(1-y)Hg+y(1-t)Bi+ytSn\}(l)$ at 600 K. Comparison with the partial ideal-solution model 653

CHAUDHARI, S. K.; KATTI, S. S. Excess molar volumes of {an *n*-alkanol (methanol, ethanol, *n*-propanol, or *n*-butanol) or an *n*-alkane (*n*-hexane, *n*-heptane, or *n*-octane) + 1,2-dichloroethane} at 298.15 K 667

Number 7, July 1989

LAVUT, E. G.; TIMOFEYEV, B. I.; YULDASHEVA, V. M. Enthalpy of formation of cobalt chloride. 673

EFIMOV, M. E.; PROKOPENKO, I. V.; MEDVEDEV, V. A.; TSIRELNIKOV, V. I.; BEREZOVSKII, G. A.; PAUKOV, I. E. Thermodynamic properties of zirconium chlorides. II. The standard molar enthalpies of formation, the low-temperature heat capacities, the standard molar entropies, and the standard molar Gibbs energies of formation of zirconium monochloride, zirconium dichloride, and zirconium tetrachloride 677

SKRIPOV, V. P.; FAIZULLIN, M. A. Thermodynamic similarity of phase-separating binary liquid mixtures having an upper critical temperature . 687

O'HARE, P. A. G.; HOPE, G. A. Standard molar enthalpy of formation at 298.15 K of the β-modification of molybdenum ditelluride 701

BAÑOS, I.; VALERO, J.; PÉREZ, P.; GRACIA, M.; GUTIÉRREZ LOSA, C. Excess molar enthalpies of mixtures containing 1,2-dichloropropane 709

LAVUT, E. G.; TIMOFEYEV, B. I.; YULDASHEVA, V. M.; RAUPOV, N. Enthalpy of formation of vanadium dichloride 715

KASPRZYCKA-GUTTMAN, T.; CHOJNACKA, I. (Vapour + liquid) equilibria of (pyridine or α-picoline + a C_6 to C_{10} *n*-alkane) at 348.15 K 721

DOHNAL, V.; PATTERSON, D. Excess molar enthalpies of {1,1,2-trichloro-trifluoroethane (freon 113) + an oxygenated solvent or a hydrocarbon} . 727

ANDERKO, A. (Vapour + liquid) equilibria of (phenol + butylbenzene or propylbenzene or isopropylbenzene or 1,2,4-trimethylbenzene or ethylbenzene) at 393.15 and 403.15 K 731

NÚÑEZ, L.; BARRAL, L.; GÓMEZ-ORELLANA, I.; RAMIL-RIO, P. E.; PAZ-ANDRADE, M. I. Excess molar enthalpies at 298.15 K of (ethyl acetate + *n*-butylmethylamine or di-*n*-propylamine or di-*n*-butylamine) and of (ethyl propionate + *n*-butylmethylamine or di-*n*-propylamine or di-*n*-butylamine) 739

ROBIE, R. A.; HASELTON, H. T., JR.; HEMINGWAY, B. S. Heat capacities and entropies at 298.15 K of $MgTiO_3$ (geikielite), ZnO (zincite), and $ZnCO_3$ (smithsonite) 743

YESAVAGE, V. F.; JOYCE, T. P. Enthalpy of 1-methylnaphthalene between 291 and 669 K at pressures to 3240 kPa 751

JIMÉNEZ, P.; ROUX, M. V.; TURRIÓN, C. Thermochemical properties of *N*-heterocyclic compounds. II. Enthalpies of combustion, vapour pressures, enthalpies of sublimation, and enthalpies of formation of 1,2,4-triazole and benzotriazole 759

LAVUT, E. G.; CHELOVSKAYA, N. V. Use of an electric furnace in a calorimetric bomb for combustion in oxygen. Determination of the standard molar enthalpy of formation of V_2O_5 765

LAINEZ, A.; GOPAL, P.; ZOLLWEG, J. A.; STREETT, W. B. Speed-of-sound measurements for liquid trichlorofluoromethane under pressure . . 773

KNAUTH, P.; SABBAH, R. Combustion calorimetry on milligram samples of hygroscopic solid substances with a CRMT rocking bomb calorimeter. Thermochemical study of ω-alkanediols at 298.15 K. Part II . . . 779

Number 8, August 1989

SKUDLARSKI, K.; DUDEK, J.; KAPAŁA, J. Thermodynamics of $\{x CdBr_2 + (1-x)CdI_2\}$(s) investigated by mass spectrometry 785

KSIĄŻCZAK, A. Vapour pressures of binary three-phase (solid + liquid + vapour) mixtures. III. Melting temperatures of the solid phases of *n*-eicosane 789

DIOT, M.; BONNETOT, B.; ABOUWASSIB, A. $\{(C_2H_5)_4N\}_2B_{10}H_{10}$ and $\{(C_2H_5)_4N\}_2B_{12}H_{12}$: heat capacity, thermodynamic properties from 5 to 300 K, and transitions 793

FRENCH, H. T. Excess enthalpies of (acetone + water) at 278.15, 288.15, 298.15, 308.15, 318.15, and 323.15 K 801

ZHAO, J.; BAO, J.; HU, Y. Excess molar enthalpies of (an alkanol + a carboxylic acid) at 298.15 K measured with a Picker calorimeter 811

YANES, C.; PÉREZ-TEJEDA, P.; MAESTRE, A. Excess molar enthalpies of (*cis*-9-octadecenoic acid + *n*-hexane or benzene or trichloroethane or tetrachloroethene) at 298.15 K 819

GARCÍA-SÁNCHEZ, F.; ROMERO-MARTÍNEZ, A.; TREJO, A. Vapour pressure, critical temperature, and critical pressure of dichloromethane . . . 823

O'Hare, P. A. G.; Susman, S.; Volin, K. J. Thermochemistry of germanium monoselenide, and the Ge–Se bond dissociation enthalpy . . . 827

Asano, M.; Kou, T. Thermochemical properties of $BaBO_2(g)$ and $Ba_3B_2O_6(s)$. 837

Bugajewski, Z.; Bylicki, A. (Solid + liquid) phase equilibria for (*N*-methylpyrrolidin-2-one + *o*-cresol 847

Connolly, J. F.; Kandalic, G. A. Thermodynamic properties of solutions of hydrogen in *n*-octane 851

Abboud, J.-L. M.; Jiménez, P.; Roux, M. V.; Turrión, C.; López-Mardomingo, C. Structural effects on the thermochemical properties of carbonyl compounds. I. Enthalpies of combustion, vapour pressures and enthalpies of sublimation, and enthalpies of formation of 2-methylpropanamide, 2,2-dimethylpropanamide, and 1-adamantyl carboxamide 859

Ewing, M. B.; Goodwin, A. R. H.; Trusler, J. P. M. Thermophysical properties of alkanes from speeds of sound determined using a spherical resonator 3. *n*-Pentane 867

Ely, J. F.; Haynes, W. M.; Bain, B. C. Isochoric (p,V_m, T) measurements on CO_2 and on ($0.982CO_2$ + $0.018N_2$) from 250 to 330 K at pressures to 35 MPa 879

NOTE

Hagopian, S.; Whittenburg, S. L. Hypersonic speed and dynamic isentropic compressibility in pyridazine 895

Number 9, September 1989

Sharma, S. C.; Joshi, I. M.; Singh, J. Excess molar volumes of (oxane + trichloromethane or tetrachloromethane or 1,2-dichloroethane or 1,1,2-trichloroethane or 1,1,2,2-tetrachloroethane or pentachloroethane) at 303.15 K 897

Murphy, K. P.; Gill, S. J. Thermodynamics of dissolution of solid cyclic dipeptides containing hydrophobic side groups 903

Niesen, V. G. (Vapor + liquid) equilibria and coexisting densities of (carbon dioxide + *n*-butane) at 311 to 395 K 915

Ghogomu, P. M.; Dellacherie, J.; Balesdent, D. Solubility of normal paraffin hydrocarbons (C_{20} to C_{24}) and some of their binary mixtures ($C_{22}+C_{24}$) and ($C_{23}+C_{24}$) in ethylbenzene 925

FILIPPOV, V. K.; KALINKIN, A. M.; VASIN, S. K. Thermodynamics of phase equilibria of aqueous (lithium sulfate + alkali-metal sulfate) (alkali metal: Na, K, and Rb), and (sodium sulfate + rubidium sulfate), at 298.15 K using Pitzer's model 935

MORIYOSHI, T.; SAKAMOTO, T.; UOSAKI, Y. (Liquid + liquid) equilibria of (water + ethanol + a C_8 alkanol) from 0.1 to 200 MPa at 298.15 K 947

NAGATA, I.; TAMURA, K. Excess molar enthalpies of (butan-2-ol + acetonitrile), (2-methylpropan-2-ol + acetonitrile or benzene), (acetonitrile + benzene), and {butan-2-ol or 2-methylpropan-2-ol + (acetonitrile + benzene)} . 955

NAGATA, I. (Liquid + liquid) equilibria of {(acetonitrile + methylcyclohexane) + methanol or ethanol or propan-1-ol or propan-2-ol or butan-1-ol or butan-2-ol} 963

MANOHAR, S.; SARADA, S.; ANANTHASWAMY, J. Thermodynamics of electrolyte solutions: an e.m.f. study of the activity coefficients of KCl in $\{KCl(m_A) + CaCl_2(m_B)\}$(aq) at 298.15, 308.15, and 318.15 K 969

CHAO, J. P.; DAI, M. Excess enthalpies of (an alkan-1-ol + tetrahydrofuran or cyclohexanone) at 298.15 K 977

YADAV, O. P. Activity coefficients and excess molar Gibbs energies of (*p*-xylene + tetrachloromethane) 985

MALLU, B. V.; NATARAJAN, G.; VISWANATH, D. S. (p, V_m, T, x) and virial coefficients of $\{xCO_2+(1-x)CO\}$ 989

DAHLMANN, U.; SCHNEIDER, G. M. (Liquid + liquid) phase equilibria and critical curves of {ethanol + dodecane or tetradecane or hexadecane or 2,2,4,4,6,8,8-heptamethylnonane)} from 0.1 MPa to 120.0 MPa. . . . 997

APELBLAT, A.; MANZUROLA, E. Solubility of ascorbic, 2-furancarboxylic, glutaric, pimelic, salicylic, and *o*-phthalic acids in water from 279.15 to 342.15 K and apparent molar volumes of ascorbic, glutaric, and pimelic acids in water at 298.15 K 1005

Number 10, October 1989

BAIDAKOV, V. G.; GURINA, T. A. An experimental study of the equation of state of superheated liquid methane 1009

LORENZANA, M. T.; LEGIDO, J. L.; JIMÉNEZ, E.; FERNÁNDEZ, J.; PÍAS, L.; ORTEGA, J.; PAZ ANDRADE, M. I. Thermodynamic properties of (a propyl ester + an *n*-alkane) at 298.15 K. I. $\{xC_2H_5CO_2C_3H_7 + (1-x)C_nH_{2n+2}\}$, ($n = 6$ to 10) 1017

ACHTERMANN, H. J.; BAEHR, H. D.; BOSE, T. K. An optical method for the determination of the (pressure, amount-of-substance density, temperature) relation. Measurements on ethene in the temperature range from 283.15 to 373.15 K at pressures up to 30 MPa 1023

GARCÍA BAONZA, V.; CÁCERES ALONSO, M., NÚÑEZ DELGADO, J. Equation of state and derived thermodynamic properties of liquid tetramethylsilane from 198 to 298 K and pressures up to 102 MPa 1045

LETCHER, T. M.; SISWANA, P. M.; WATT, P. VAN DER; RADLOFF, S. Phase equilibria for (an alkanol + *p*-xylene + water) at 298.2 K . . . 1053

SIJPKES, A. H.; ROSSUM, P. VAN; RAAD, J. S.; SOMSEN, G. Heat capacities and volumes of some polybasic carboxylic acids in water at 298.15 K . . 1061

DUSCHEK, W.; KLEINRAHM, R.; WAGNER, W.; JAESCHKE, M. Measurement of the (pressure, density, temperature) relation of Ekofisk natural gas in the temperature range from 273.15 to 323.15 K at pressures up to 8 MPa . 1069

MORSS, L. R.; HAAR, C. M.; MROCZKOWSKI, S. Standard molar enthalpy of formation of neodymium hydroxide 1079

SIPOWSKA, J. T.; GRAHAM, R. C.; NEELY, B. J.; OTT, J. B.; IZATT, R. M. Excess enthalpies for (ethane + methanol) at 298.15 and 348.15 K and 7.5 and 15 MPa, and at 323.15 K at 7.5, 10, and 15 MPa 1085

WYERS, G. P.; CORDFUNKE, E. H. P.; OUWELTJES, W. The standard molar enthalpies of formation of the lithium zirconates 1095

GALLARDO, M. A.; GARCÍA VICENTE, I.; VELASCO, I.; OTÍN, S. Excess volumes of some (1-chloroalkane or 1-bromoalkane + benzene)s at 298.15 K . . 1101

Number 11, November 1989

KIRKLIN, D. R.; CHURNEY, K. L.; DOMALSKI, E. S. Enthalpy of combustion of 1,4-dimethylcubane dicarboxylate 1105

ROGERS, D. W.; DEJROONGRUANG, K. Enthalpies of hydrogenation of the dimethylpentenes, ethylpentenes, methylethylbutene, and trimethylbutene. 1115

STEELE, W. V.; CHIRICO, R. D.; HOSSENLOPP, I. A.; NGUYEN, A.; SMITH, N. K.; GAMMON, B. E. The thermodynamic properties of 1,2,3,4- and 5,6,7,8-tetrahydroquinolines 1121

VINE, M. D.; WORMALD, C. J. The enthalpy of ethanol 1151

BAIDAKOV, V. G.; KAVERIN, A. M. Ultrasonic speed in superheated liquid oxygen 1159

CHAO, J. P.; DAI, M.; WANG, Y. X. Studies of thermodynamic properties of binary mixtures containing an alcohol. XII. Excess molar enthalpies of (an alkan-1-ol + *N*,*N*-dimethylformamide or dimethyl solfoxide) at 298.15 K 1169

PIKKARAINEN, L. Excess enthalpies of (2,2,2-trifluoroethanol + formamide or *N*-methylformamide or *N*,*N*-dimethylformamide) 1177

VISWANATHAN, R.; BALASUBRAMANIAN, R.; MATHEWS, C. K. Vaporization thermodynamics of the metal-rich molybdenum telluride Mo_3Te_4: a high-temperature mass-spectrometric study 1183

BRUMLEVE, T. R.; MUCKLEJOHN, S. A.; O'BRIEN, N. W. The preparation and vapour pressures of the indium(I) halides and the standard molar Gibbs free energy change for formation of InX from In(g) and X_2(g), (X = Cl, Br, or I). 1193

AMIGO, A.; LEGIDO, J. L.; BRAVO, R.; PAZ ANDRADE, M. I. Excess molar enthalpies of (heptan-1-ol + an *n*-alkane) at 298.15 and 308.15 K . . 1207

ALONSO, R.; CORRALES, J. A. Excess molar volumes of (methylcyclohexane + an alkanol) at 298.15 K. II. Results for 2-methylpropan-1-ol and butan-2-ol 1213

Number 12, December 1989

YANES, C.; PÉREZ-TEJEDA, P.; MAESTRE, A. Excess molar volumes and excess refractive indices of (*cis*-9-octadecenoic acid + *n*-hexane or cyclohexane or benzene or trichloroethene or tetrachloroethene) at 298.15 K . . . 1217

JUNQUERA, E.; AICART, E.; TARDAJOS, G. Speed of sound and isentropic compressibility of (1-chlorobutane + *n*-undecane or *n*-dodecane or *n*-tetradecane or *n*-hexadecane) at 298.15 K 1223

KSIĄŻCZAK, A. Vapour pressures of binary three-phase (solid + liquid + vapour) mixtures. IV. Melting temperatures of the solid phases of *n*-octadecane and of *n*-nonadecane 1231

KSIĄŻCZAK, A. Vapour pressures of binary three-phase (solid + liquid + vapour) mixtures. V. (Phenol + benzene), (*p*-methylphenol + benzene), and (*p*-chlorophenol + benzene) 1237

KALRA, K. C.; SINGH, K. C.; SPAH, D. C. Ultrasonic speeds and isentropic compressibilities of (1,2-dibromoethane + an aromatic hydrocarbon) at 298.15 K 1243

HANDA, Y. P.; YAMAMURO, O.; OGUNI, M.; SUGA, H. Low-temperature heat capacities of xenon and krypton clathrate hydrates 1249

TOSCANI, S.; FIGUIÈRE, P.; SZWARC, H. A magnetic-suspension apparatus to measure densities of liquids as a function of temperature at pressures up to 100 MPa. Application to *n*-heptane 1263

ESTESO, M. A.; FERNÁNDEZ-MÉRIDA, L.; GRANDOSO, D. M. Ionic interactions in aqueous solutions of organic electrolytes. Activity coefficients for (sodium chloride + sodium hydrogen malonate)(aq) at 298.15 K. Applications of Scatchard's and of Pitzer's equations 1279

MCELROY, P. J.; BATTINO, R.; DOWD, M. K. Compression-factor measurements on methane, carbon dioxide, and (methane + carbon dioxide) using a weighing method 1287

SIJPKES, A. H.; SOMSEN, G. Enthalpies of dilution of benzamide, cyclohexanecarboxamide, and *N*-acetyl-*N'*-methyl-L-cyclohexylalaninamide dissolved in *N*,*N*-dimethylformamide at 298.15 K 1301

CHIRICO, R. D.; KNIPMEYER, S. E.; NGUYEN, A.; STEELE, W. V. The thermodynamic properties of biphenyl 1307

INDEX TO VOLUME 21 1333